AD)				

Award Number: DAMD17-01-1-0116

TITLE: Thromboxane Synthase and Prostate Cancer Progression

PRINCIPAL INVESTIGATOR: Kenneth V. Honn, Ph.D.

David Grignon, M.D. Daotai Nie, Ph.D.

Graham Pidgeon, Ph.D.

CONTRACTING ORGANIZATION: Wayne State University

Detroit, MI 48202

REPORT DATE: July 2005

TYPE OF REPORT: Final Addendum

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE	2. REPORT TYPE	3. DATES COVERED
01-07-2005	Final Addendum	15 Jun 2001 – 14 Jun 2005
4. TITLE AND SUBTITLE	Tinal Addendant	5a. CONTRACT NUMBER
Thromboxane Synthase and F	Prostate Cancer Progression	56. GRANT NUMBER DAMD17-01-1-0116
		5c. PROGRAM ELEMENT NUMBER
6. AUTHOR(S) Kenneth V. Honn, Ph.D.		5d. PROJECT NUMBER
David Grignon, M.D. Daotai Nie, Ph.D.		5e. TASK NUMBER
Graham Pidgeon, Ph.D.		5f. WORK UNIT NUMBER
7. PERFORMING ORGANIZATION N Wayne State University Detroit, MI 48202	NAME(S) AND ADDRESS(ES)	8. PERFORMING ORGANIZATION REPORT NUMBER
9. SPONSORING / MONITORING AC U.S. Army Medical Research : Fort Detrick, Maryland 21702		10. SPONSOR/MONITOR'S ACRONYM(S)
1 of Bothon, Walyland 21702		11. SPONSOR/MONITOR'S REPORT NUMBER(S)
42 DICTRIBUTION / AVAIL ADILITY	CTATEMENT	

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for Public Release; Distribution Unlimited

13. SUPPLEMENTARY NOTES

Original contains color plates ALL DTIC reproductions will be in black and white

14. ABSTRACT

The initiation and progression of prostate cancer remain not well understood to enable development of interventional therapy. Thromboxane synthase is an enzyme downstream of cyclooxygenase, utilizing prostaglandin H to form thromboxane A2. Using immunohistochemistry analysis, we found that 25% of clinical prostate tumor specimens had strong expression of thromboxane synthase; 33% of cases had medium expression and 42% of cases had weak expression of thromboxane synthase. Prostate cancer cells isolated from lymph node metastasis had higher levels of thromboxane synthase expression and activity than those isolated from the primary tumor sites in an animal model. We cloned and sequenced full-length thromboxane synthase cDNA from PC-3 cells and constructed an expression vector. Increased expression of thromboxane synthase in DU145 cells was found to stimulate cell migration. Inhibition of thromboxane synthase or blockade of thromboxane A2 function inhibited prostate cancer cell migration. Further we found that PCa cells express receptors for TXA2 and stimulation with TXA2 mimetic led to the activation of RhoA and cell retraction. Our study suggest that thomboxane synthase and its eicosanoid product play a contributory role in prostate cancer progression.

15. SUBJECT TERMS

Prostate cancer, antiogenesis, eicosanoids, platelets, thromboxanes

16. SECURITY CLASSIFICATION OF:		17. LIMITATION	18. NUMBER	19a. NAME OF RESPONSIBLE PERSON		
		OF ABSTRACT	OF PAGES	USAMRMC		
a. REPORT U	b. ABSTRACT	c. THIS PAGE U	טט	11	19b. TELEPHONE NUMBER (include area code)	

TABLE OF CONTENTS

Cover	
Sf 298 2)
Introduction2	1
Body 5	
List of technical objectives	
Key Research Accomplishments 8	
Reportable Outcomes9)
Conclusions)

INTRODUCTION

The hemostatic system plays an important role in the regulation of tumor angiogenesis and progression. Prostate cancer is one of most common cancers affecting American men. The initiation and progression of prostate cancer remain not well understood to enable rational development of interventional therapy. In this proposal, we propose to study thromboxane synthase and its product, thromboxane A₂, in prostate cancer progression. Thromboxane synthase is an enzyme downstream cyclooxygenase, utilizing prostaglandin H to form thromboxane A2. Thromboxane A2 is a potent inducer of platelet aggregation which can subsequently lead to coagulation and thrombosis, a hematological complication affecting about 15% to 20% of cancer patients and the second leading cause of death in cancer patients. Platelet aggregation also release a plethora of angiogenesis regulators and fibrin deposition to facilitate angiogenesis and formation of tumor stroma. In addition, it has been demonstrated that thromboxane A2 also can directly modulate endothelial cell angiogenic responses. It is our working hypothesis that TXA2 may play an important role in prostate tumor progression and that this functional role of TXA2 is achieved by modulating tumor cell motility, endothelial angiogenic responses, and platelet aggregation. TXA2 produced by PCa cells can promote PCa cell migration as an autocoid and modulate endothelial cell angiogenic responses and stimulate platelet aggregation as a paracrine factor. Platelet aggregation release various angiogenesis regulators and cause coagulation, which subsequently leads to deposition of fibrin at tumor sites. These events, collectively, promote tumor angiogenesis and growth. The proposed work will provide significant insights into how prostate cancer cells regulate cell migration and hemostatic system to facilitate tumor angiogenesis, growth, and metastasis. The knowledge obtained from the proposed work will identify key targets (TX synthase and TXA₂ receptor) to develop interventional therapy for prostate cancer, advancing the program's eventual goal to eliminate prostate cancer.

BODY OF REPORT

List of Technical Objectives

Modified Task 2. Study the effect of increased expression of wild type TX synthase in DU145 cells on cell migration, proliferation, and apoptosis.

- a. Convert the mutated TX synthase expression construct to wild-type, through site directed mutagenesis approach.
- b. Transfect Du145 cells using the new, wild-type TX synthase expression construct and study the biosynthesis of TXA2.
- c. Plate the transfected DU145 cells and vector control cells and study the proliferation rate using MTS proliferation assay kit.
- d. Plate TX synthase transfected DU145 cells, and their vector control, in 10 cm dishes and study whether TX synthase overexpression confers on DU145 cells resistance to apoptosis induced by COX inhibitor NS398, as assessed by DNA laddering and apoptosis induced by COX inhibitor NS398, as assessed by DNA laddering and apoptosis ELISA for DNA fragmentation.
- e. Plate TX synthase transfected DU145 cells and vector control cells in fibronectin-precoated Boyden chambers for migration assay as a standard procedure.

Research Progress

When we submitted our request for a one year no-cost extension we listed Modified Task 2 above as our stated objective. However shortly after we receiving the extension we initiated a collaboration with Dr. Anthony Ashton of the Albert Einstein Medical School. Dr. Ashton had developed antibodies individually recognizing the two thromboxane receptors, i.e. $TP\alpha$ and $TP\beta$. These antibodies which were useful in immunohistochemistry gave us the unique opportunity to conduct a clinical study on expression of $TP\alpha$ and $TP\beta$ in human prostate tissues. Therefore we modified our objectives to the analysis of $TP\alpha$ and $TP\beta$ expression. The results of which are presented below.

Expression of TP-beta and TP-alpha in human prostate

Expression of TP-beta and TP-alpha was analyzed by immunohistochemistry in 51 radical prostatectomy specimens, which included 25 organ-confined (pT2) and 26 locally advanced diseases (16 pT3a and 10 pT3b). Of those 25 organ-confined prostate cancers, 10, 14 and 1 had a Gleason score 6, 7 and 8 respectively, whereas of those 26 locally advanced diseases, 21, 4 and 1 had a Gleason score 7, 8 and 9 respectively. Immunoreactivity was evaluated semiquantitatively according to staining intensity and percentage of cells stained. A staining score, which was determined by the percentage of tumor cells with moderate and strong staining intensity, was assigned to each tumor. Mann-Whitney test was used for statistical analysis and a p value of <0.05 was considered significant.

In normal prostatic glands, TP-beta was mainly expressed in the basal cells. Increased expression in luminal cells was found in areas of atrophy, basal cell hyperplasia, inflammation and transitional cell metaplasia. Significantly increased TP-beta expression was also found in high-grade prostatic intraepithelial neoplasia (Figure 1). The expression of TP-alpha was very focal and week in non-neoplastic prostate. In high-grade prostatic intraepithelial neoplasia, the expression of TP-alpha was variably increased (Figure 2). Both markers were heterogeneously expressed in prostatic adenocarcinomas. When the cases was separated into 2 groups according to pathologic stage (pT), a tendency of increased expression of both markers was noted in the advanced tumors (Table 1). However, these differences were not statistically significant. Interestingly, when the tumors were categorized according to Gleason score, expression of both markers was significantly increased in moderately to poorly differentiated tumors (Gleason score ≥7) compared to moderately well differentiated tumors (Gleason score 6) (Table 2, Figure 1 and 2). There was no further significant differences

in the expression of either marker between moderately poorly differentiated tumors (Gleason score 7) and poorly differentiated tumors (Gleason score 8 and higher). These results suggest that TP-alpha and TP-beta may play important roles in the initiation and early progression of prostate cancer.




Figure 1. TP-beta expression in human prostate.

A, normal; B, high-grade prostatic intraepithelial neoplasia; C, Gleason score 6 prostatic adenocarcinoma; and D, Gleason score 7 prostatic adenocarcinoma.




Figure 2. TP-alpha expression in human prostate.

A, normal; B, high-grade prostatic intraepithelial neoplasia; C, Gleason score 6 prostatic adenocarcinoma; and D, Gleason score 7 prostatic adenocarcinoma.

Table 1. TP expression in prostate according to pathologic stage

Pathologic Stage	N	TP-BETA	TP-ALPHA
		Median	Median
		(range)	(range)
pT2	25	50	20
		(5-100)	(5-70)
рТ3	26	70	30
		(5-100)	(5-80)
P Value	-	0.1171	0.2443

TABLE 2. TP EXPRESSION IN PROSTATE ACCORDING TO GLEASON SCORE

Median (range)	Median (range)
30	15
(5-80)	(5-70)
70	30
(5-100)	(5-80)
0.0082	0.0109
	70 (5-100)

Key Research Accomplishments

We were able to demonstrate that antibodies individually recognizing $TP\alpha$ and $TP\beta$ were suitable for immunohistochemical studies on human prostate tissue. Further we were able to show a significant increase in expression of $TP\alpha$ and $TP\beta$ with increased Gleason grade.

Reportable Outcomes

• Presentations

Thromboxane Synthase and TP Receptors In Tumor Progression

Kenneth V. Honn

9th International Conference on Eicosanoids and Other Bioactive Lipids in Cancer, Inflammation, and Related Diseases. San Francisco, CA. September 11-14, 2005. (Podium Presentation)

Abstracts Published

Thromboxane Synthase and TP Receptors In Tumor Progression

Kenneth V. Honn

9th International Conference on Eicosanoids and Other Bioactive Lipids in Cancer, Inflammation, and Related Diseases. San Francisco, CA. September 11-14, 2005.

Thromboxane Synthase in Human Breast Cancer Cells

Yande Guo, Keqin Tang, Yinlong Cai, Yan Qiao, Daotai Nie, and Kenneth V. Honn 9th International Conference on Eicosanoids and Other Bioactive Lipids in Cancer, Inflammation, and Related Diseases. San Francisco, CA. September 11-14, 2005.

- Patents applied: None
- Degress obtained that are supported by this award: None
- Development of cell lines, tissue or serum repositories: None
- Funding applied or obtained: Yes

Based on the above described findings as well other findings previous reported we submitted an application to the National Institutes of Health entitled "Role of Thromboxane in Prostate Cancer Progression" this grant received a sufficient priority score and will be funded next month.

Conclusions:

Our studies over the past one year extension of this grant demonstrated that thromboxane receptors $TP\alpha$ and $TP\beta$ were both expressed in human prostate tissues. The expression was weak in the non-neoplastic prostate. Expression was found in high grade prostatic interepithelial neoplasia and both markers increased expression in high Gleason grade tumors verses low Gleason tumors.




Figure 2